

Cataract surgery requires eye drops before and after surgery. These are necessary to assist in the prevention of inflammation and infection to achieve your best visual outcome. We have come up with a simplified solution, incorporating all 3 drops into just 1 bottle of drops. You will purchase this bottle of drops at the time of scheduling your surgery.

- The Imprimis eye drops are \$55 a bottle.
- One bottle should be enough medicine for 1 eye surgery, if using as directed.
- You will need to purchase a 2nd bottle of drops prior to your 2nd eye surgery.
- It is not available through the pharmacy or your insurance.
- You can use your flex pay or HSA to pay for this medicine.



You will start the drops in the surgical eye 3 days prior to surgery, 1 drop 4 times a day.

*The day of surgery, you can take your drop first thing in the morning and then follow your doctors instructions for the remainder of Day 1.

The first week, you take one drop 4 times a day.

The second week, you take one drop 3 times a day.

The third week, you take one drop 2 times a day.

The fourth (final) week, you take one drop once a day.

Tips for Proper Eye Drop Administration:

***Shake the bottle of drops well before using!**

***Gently tilt the bottle to instill the drops. If you need to squeeze the bottle, do this very gently as the bottle does not require much pressure to dispense a drop. This will ensure you get the appropriate number of drops from the bottle.**

The chart on the back of this sheet will help you keep track of when to take your drops.

3 days prior to surgery (Date): _____ Start your drops, 1 drop 4 times a day in ☐ Right ☐ Left eye. Shake the bottle of drops well before using.

After surgery	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Week 1	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐
_____	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐
	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐
Week 2	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐
_____	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐
Week 3	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐

Week 4	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start							

3 days prior to surgery (Date): _____ Start your drops, 1 drop 4 times a day in ☐ Right ☐ Left eye. Shake the bottle of drops well before using.

After surgery	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Week 1	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐
_____	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐
	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐	4 ☐
Week 2	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐
_____	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐	3 ☐
Week 3	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐	2 ☐

Week 4	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐	1 ☐
Start							

Please review the tips of proper eye drop administration highlighted on the back of this sheet.